## **ASTHMA CARE PLAN FOR EDUCATION** AND CARE SERVICES

CONFIDENTIAL: Staff are trained in Asthma First Aid (see overleaf) and can provide routine asthma medication as authorised in this care plan by the treating doctor. Please advise staff in writing of any changes to this plan.

To be completed by the treating doctor and parent/guardian, for supervising staff and emergency medical personnel.

PLEASE PRINT CLEARLY Plan date /\_\_\_\_/20\_ DOB: \_ Student's name: \_ Review date \_/\_\_\_/20\_

#### MANAGING AN ASTHMA ATTACK

Staff are trained in Asthma First Aid (see overleaf). Please write down anything different this student might need if they have an asthma attack:

#### **DAILY ASTHMA MANAGEMENT**

This student's usual asthma signs:	Frequency and severity:		Known triggers for this student's asthma (e.g. exercise*, colds/flu, smoke) —	
Cough	Daily/most days		please detail:	
Wheeze	Frequently (more than 5 x per	Frequently (more than 5 x per year)		
Difficulty breathing	Occasionally (less than 5 x per	Occasionally (less than 5 x per year)		
Other (please describe):	Other (please describe)			
Does this student usually tell an adult if s/he is having trouble breathing?		Yes	No	
Does this student need help to take asthma medication?		Yes	No	
Does this student use a mask with a spacer?			No	
*Does this student need a blue/grey reliever puffer medication before exercise?			No	

#### **MEDICATION PLAN**

NAME OF MEDICATION AND COLOUR

If this student needs asthma medication, please detail below and make sure the medication and spacer/mask are supplied to staff.

**DOSE/NUMBER OF PUFFS** 

DOCTOR	PAREN	PARENT/GUARDIAN		EMERGENCY CONTACT INFORMATION	
Name of doctor	attachmen	I have read, understood and agreed with this care plan and any attachments listed. I approve the release of this information to staff and emergency medical personnel. I will notify the staff in writing if there are any changes to these instructions. I understand staff will seek emergency medical help as needed and that I am responsible for payment of any emergency medical costs.		Contact name	
Address	writing if the staff will			Phone	
Phone	Signatu	re Date		Mobile	
Cignoturo	Data Nama			Email	

TIME REQUIRED

PHOTO OF STUDENT

# **ASTHMA FIRST AID**

## **Blue/Grey Reliever**

Airomir, Asmol, Ventolin or Zempreon and Bricanyl

Blue/grey reliever medication is unlikely to harm, even if the person does not have asthma



DIAL TRIPLE ZERO (000) FOR AN AMBULANCE IMMEDIATELY IF THE PERSON:

- is not breathing
- suddenly becomes worse or is not improving
- is having an asthma attack and a reliever is not available
- is unsure if it is asthma
- has a known allergy to food, insects or medication and has SUDDEN BREATHING DIFFICULTY, GIVE ADRENALINE AUTOINJECTOR FIRST (if available)





SIT THE PERSON UPRIGHT

- Be calm and reassuring
- Do not leave them alone

2



GIVE 4 SEPARATE PUFFS OF RELIEVER PUFFER

- Shake puffer
- Put 1 puff into spacer
- Take 4 breaths from spacer
  - Repeat until 4 separate puffs have been taken



If using **Bricanyl** (5 years or older)

- Do not shake. Open, twist around and back, and take a deep breath in
- Repeat until 2 separate inhalations have been taken

If you don't have a spacer handy in an emergency, take  $\underline{1}$  puff as you take  $\underline{1}$  slow, deep breath and hold breath for as long as comfortable. Repeat until all puffs are given

3



WAIT 4
MINUTES

If breathing does not return to normal, give
 4 more separate puffs of reliever as above



Bricanyl: Give 1 more inhalation

### IF BREATHING DOES NOT RETURN TO NORMAL





DIAL TRIPLE ZERO (000)

- Say 'ambulance' and that someone is having an asthma attack
- Keep giving <u>4 separate puffs every</u>
   <u>4 minutes</u> until emergency assistance arrives



**Bricanyl:** Give 1 more inhalation <u>every 4 minutes</u> until emergency assistance arrives







